



Mindfulness Meditation and Evidence of Brain Changes: A Review of the Literature


Candice Dexter
Donna Oropall

UB Psychology/Human Services Department
University of Bridgeport, Bridgeport, CT





Abstract

Mindfulness meditation is a practice that can develop greater awareness of the mind and body. It has been used to treat stress, anxiety, chronic pain (12) and depression. Current studies shed light on research findings associated with the effects of mindfulness meditation on the brain. This literature review will reveal evidence of neuroplasticity through functional imaging, such as the fMRI. Likewise, evidence of emotional regulation involving amygdala activation has been presented in some studies (4,2). Additionally, increased gray matter density in the brain stem has been documented after meditation therapy (MT) using magnetic resonance imaging(9). Researchers have additionally uncovered significant improvements in immune system function among meditators (10).



Main Arguments

There are a variety of benefits resulting from mindfulness meditation that include stress and anxiety reduction, emotional regulation and management of chronic pain. In recent years, science has been catching up and confirming many of these effects empirically as researchers across disciplines investigate the neurological and physiological effects of mindful meditation (9). Meditation had been used to help treat cancer patients manage stress (13). There is evidence of neuroplastic effects in the brain after meditation training (MT) resulting in improved self-regulation, cognition and neural processing (1,12). Most research makes use of fMRI imaging that compares the brains of study participants before and after mindfulness meditation (14). The effects of such programs have demonstrated numerous neural changes (6). For instance amygdala activity after MT suggests potential for lasting changes in brain function, specifically emotional responses (2,3,5). Right side amygdala activity decreased after a mindfulness based stress reduction program, suggesting reduced reaction to negative emotional stimulus in social anxiety disorder (SAD) patients (2). Amygdala connectivity changes were similarly correlated with the Beck Anxiety Inventory (4) in general anxiety disorder (GAD) patients following MBSR, along with reduced anxiety and stress symptoms (7) .




Key Questions

- To examine the association between mindfulness meditation and the neuroplasticity of the brain.
- To determine the relationship between MT and reduction in stress, depression and anxiety.

Conclusion

Mindfulness meditation has been shown to change the way people think, the way we respond to stimuli and even the physical structure of our brains. A more comprehensive understanding of the physical and neural changes that MT causes, and the capacity for the brain to change, could be used to improve upon current MT therapies and their application. Similarly, MT related changes in neuro-activity could lead to new psychological treatments in coming years. For instance better understanding of the plasticity of the amygdala could be helpful in developing new treatments for mood disorders, fears, addiction and anxiety. MT's mind, body, and internal awareness benefits could likewise be applied to treatment of borderline personality disorder. Channeling of attention, and control of focus could aid people with ADD. MBSR therapeutic programs have been adopted for use in mainstream medicine and used as templates for other psychotherapies, and should continue to be built upon. Additionally, study of other stress reduction techniques, such as meditative prayer (15) along with mindfulness meditation should be studied in parallel to potentially expand treatment options based on the most effective technique for a given patient.



References

1-Allen, M., Dietz, M., Blair, K. S., van Beek, M., Rees, G., Vestergaard-Poulsen, P., ... & Roepstorff, A. (2012). Cognitive-affective neural plasticity following active-controlled mindfulness intervention. *The Journal of Neuroscience*, 32(44), 15601-15610

10-Davidson, R. J., Kabat-Zinn, J., Schumacher, J., Rosenkranz, M., Muller, D., Santorelli, S. F., & ... Sheridan, J. F. (2003). Alterations in brain and immune function produced by mindfulness meditation. *Psychosomatic Medicine*, 65(4), 564-570. doi:10.1097/01.PSY.0000077505.67574.E

3-Desbordes G, Negi LT, Pace TW, Wallace BA, Raision CL, Schwartz EL. Effects of mindfulness and compassion meditation training on amygdala response to emotional stimuli in an ordinary, non-meditative state. *Front Hum Neurosci*. 2012;6:292.

6-Farb N., Segal Z. V., Mayberg H., Bean J., McKeon D., Fatima Z., & Anderson A. K. (2007). Attending to the present: mindfulness meditation reveals distinct neural modes of self-reference. *Social Cognitive and Affective Neuroscience*, 2(4), 313-322. doi:10.1093/scan/nsm030

2-Goldin PR, Gross JJ. Effects of mindfulness-based stress reduction (MBSR) on emotion regulation in social anxiety disorder. *Emotion*. 2010;10:83-91.

4-Hölzel BK, Hoge EA, Greve DN, Gard T, Creswell JD, Brown KW, Barrett LF, Schwartz C, Vaitl D, Lazar SW. Neural mechanisms of symptom improvements in generalized anxiety disorder following mindfulness training. *Neuroimage Clin*. 2013;2:448-458

12- Hölzel B., Lazar S., Gard T., Schuman-Olivier Z., Vago, D., Ott U. (2011). How does mindfulness meditation work? Proposing mechanisms of action from a conceptual and neural perspective. *Perspectives on Psychological Science*, 6(6), 537-559

7-Kabat-Zinn J., Massion A., Kristeller J., Peterson L., Fletcher K., Pbert L.,... et al. Effectiveness of a meditation-based stress reduction program in the treatment of anxiety disorders. *American Journal of Psychiatry*. 1992;149:936-43.

14 Kabat-Zinn J., & University of Massachusetts Medical Center/Worcester. (1991). Full catastrophe living: Using the wisdom of your body and mind to face stress, pain, and illness. New York, N.Y: Pub. by Dell Pub., a division of Bantam Doubleday Dell Pub. Group

11- Luders E., Toga, A. W., Lepore N., & Gaser C. (2009). The underlying anatomical correlates of long-term meditation: larger hippocampal and frontal volumes of gray matter. *Neuroimage*, 45(3), 672-678.

15-Newberg A., Pourdehnad M., Alavi A., d'Aquili E (2003). CEREBRAL BLOOD FLOW DURING MEDITATIVE PRAYER: PRELIMINARY FINDINGS AND METHODOLOGICAL ISSUES. *Perceptual and Motor Skills*, 97(2).

13-Speca M., Carlson L. E., Goodey E., & Angen M. (2000). A randomized, wait-list controlled clinical trial: the effect of a mindfulness meditation-based stress reduction program on mood and symptoms of stress in cancer outpatients. *Psychosomatic medicine*, 62(5), 613-622.

5- Taylor V. A., Grant J., Daneault, V., Scavone G., Breton E., Roffe-Vidal S., ... Beauregard M. (2011). Impact of mindfulness on the neural responses to emotional pictures in 9-Gray matter

Vestergaard-Poulsen P., van Beek M., Skewes J., Bjarkam C. R., Stubberup M., Bertelsen J., & Roepstorff A. (2009). Long-term meditation is associated with increased gray matter density in the brain stem. *Neuroreport*, 20(2), 170-174.




Figure 5. Right dorsal amygdala blood oxygenation level-dependent (BOLD) contrast signal time series during reacting to negative self-beliefs and breath-focused attention in social phobics (SP) at both pre- and post-mindfulness-based stress reduction (MBSR). * $p < .05$. Rate = negative emotion rating; React = reacting to the negative self-belief; Breath-Focus = instruction to focus attention on breath sensation.

Other research indicates improved immune system function among MBSR participants who show elevated levels of antibody titers compared with non-meditators (10). Long-term meditators have shown an increase in gray matter density(9,11).